

The Forty-Seventh Annual William Lowell Putnam Competition
Saturday, December 6, 1986

A-1 Find, with explanation, the maximum value of $f(x) = x^3 - 3x$ on the set of all real numbers x satisfying $x^4 + 36 \leq 13x^2$.

A-2 What is the units (i.e., rightmost) digit of

$$\left\lfloor \frac{10^{20000}}{10^{100} + 3} \right\rfloor?$$

A-3 Evaluate $\sum_{n=0}^{\infty} \operatorname{Arccot}(n^2 + n + 1)$, where $\operatorname{Arccot} t$ for $t \geq 0$ denotes the number θ in the interval $0 < \theta \leq \pi/2$ with $\cot \theta = t$.

A-4 A transversal of an $n \times n$ matrix A consists of n entries of A , no two in the same row or column. Let $f(n)$ be the number of $n \times n$ matrices A satisfying the following two conditions:

- (a) Each entry $\alpha_{i,j}$ of A is in the set $\{-1, 0, 1\}$.
- (b) The sum of the n entries of a transversal is the same for all transversals of A .

An example of such a matrix A is

$$A = \begin{pmatrix} -1 & 0 & -1 \\ 0 & 1 & 0 \\ 0 & 1 & 0 \end{pmatrix}.$$

Determine with proof a formula for $f(n)$ of the form

$$f(n) = a_1 b_1^n + a_2 b_2^n + a_3 b_3^n + a_4,$$

where the a_i 's and b_i 's are rational numbers.

A-5 Suppose $f_1(x), f_2(x), \dots, f_n(x)$ are functions of n real variables $x = (x_1, \dots, x_n)$ with continuous second-order partial derivatives everywhere on \mathbb{R}^n . Suppose further that there are constants c_{ij} such that

$$\frac{\partial f_i}{\partial x_j} - \frac{\partial f_j}{\partial x_i} = c_{ij}$$

for all i and j , $1 \leq i \leq n$, $1 \leq j \leq n$. Prove that there is a function $g(x)$ on \mathbb{R}^n such that $f_i + \partial g / \partial x_i$ is linear for all i , $1 \leq i \leq n$. (A linear function is one of the form

$$a_0 + a_1 x_1 + a_2 x_2 + \dots + a_n x_n.)$$

A-6 Let a_1, a_2, \dots, a_n be real numbers, and let b_1, b_2, \dots, b_n be distinct positive integers. Suppose that there is a polynomial $f(x)$ satisfying the identity

$$(1-x)^n f(x) = 1 + \sum_{i=1}^n a_i x^{b_i}.$$

Find a simple expression (not involving any sums) for $f(1)$ in terms of b_1, b_2, \dots, b_n and n (but independent of a_1, a_2, \dots, a_n).

B-1 Inscribe a rectangle of base b and height h in a circle of radius one, and inscribe an isosceles triangle in the region of the circle cut off by one base of the rectangle (with that side as the base of the triangle). For what value of h do the rectangle and triangle have the same area?

B-2 Prove that there are only a finite number of possibilities for the ordered triple $T = (x - y, y - z, z - x)$, where x, y, z are complex numbers satisfying the simultaneous equations

$$x(x-1) + 2yz = y(y-1) + 2zx = z(z-1) + 2xy,$$

and list all such triples T .

B-3 Let Γ consist of all polynomials in x with integer coefficients. For f and g in Γ and m a positive integer, let $f \equiv g \pmod{m}$ mean that every coefficient of $f - g$ is an integral multiple of m . Let n and p be positive integers with p prime. Given that f, g, h, r and s are in Γ with $rf + sg \equiv 1 \pmod{p}$ and $fg \equiv h \pmod{p}$, prove that there exist F and G in Γ with $F \equiv f \pmod{p}$, $G \equiv g \pmod{p}$, and $FG \equiv h \pmod{p^n}$.

B-4 For a positive real number r , let $G(r)$ be the minimum value of $|r - \sqrt{m^2 + 2n^2}|$ for all integers m and n . Prove or disprove the assertion that $\lim_{r \rightarrow \infty} G(r)$ exists and equals 0.

B-5 Let $f(x, y, z) = x^2 + y^2 + z^2 + xyz$. Let $p(x, y, z), q(x, y, z), r(x, y, z)$ be polynomials with real coefficients satisfying

$$f(p(x, y, z), q(x, y, z), r(x, y, z)) = f(x, y, z).$$

Prove or disprove the assertion that the sequence p, q, r consists of some permutation of $\pm x, \pm y, \pm z$, where the number of minus signs is 0 or 2.

B-6 Suppose A, B, C, D are $n \times n$ matrices with entries in a field F , satisfying the conditions that AB^T and CD^T are symmetric and $AD^T - BC^T = I$. Here I is the $n \times n$ identity matrix, and if M is an $n \times n$ matrix, M^T is its transpose. Prove that $A^T D + C^T B = I$.